

PM/97-10

May 15, 1997

ANALYTIC CONSTRAINTS FROM EW SYMMETRY BREAKING IN THE MSSM¹

Gilbert MOULTAKA

*Physique Mathématique et Théorique
UPRES-CNRS A 5032,*

Université Montpellier II, F34095 Montpellier Cedex 5, France

(Talk given at XXXIInd Rencontres de Moriond- 15-22 March 1997,
ELECTROWEAK INTERACTIONS AND UNIFIED THEORIES)

Abstract

We report on how a straightforward (albeit technically involved) analytic study of the 1-loop effective potential in the Minimal Supersymmetric Standard Model (MSSM), modifies the usual electroweak symmetry breaking (EWSB) conditions involving $\tan\beta$ and the other free parameters of the model. The study implies new constraints which (in contrast with the existing ones like $1 \leq \tan\beta \leq m_t/m_b$) are fully model-independent and exclude more restrictively a region around $\tan\beta \sim 1$. Further results of this study will be only touched upon here.

(to appear in the proceedings)

¹supported in part by EC contract CHRX-CT94-0579

Determining the structure of the true vacuum is a central task in particle physics. It encodes the mass spectrum, the way symmetries are broken and elementary fields interact. Supersymmetric theories give specific properties to the vacuum structure, some of which survive or remain approximately true after the soft breaking of supersymmetry. An attractive feature of this symmetry breaking is that it can trigger (together with a heavy top quark), the spontaneous breaking of the electroweak symmetry [1]. However, such a phenomenon should be accompanied by some extra constraints to ensure that neither charge/color breaking nor unbounded from below potential occur [2]. In the absence of any underlying mechanism that dictates those constraints, one usually resorts to a phenomenological determination using the approximation at hand of the true effective potential of the theory. It is then useful to keep in mind some essential features:

i) The true effective potential V_{eff} should be renormalization scale invariant. *ii)* The only gauge parameter independent and physically relevant values of V_{eff} are those at stationary points. *iii)* the renormalization group (RG) improved effective potential $\bar{V}_{eff}$ can miss large logarithms when there are more than one (mass) scale in the theory, as is the case in the MSSM. If not properly treated, the scale at which leading logs are actually resummed becomes fuzzy and with it the approximate scale invariance of $\bar{V}_{eff}$. *iv)* A local minimum is characterized by both 1st and 2nd order derivatives with respect to *all* scalar fields in the theory. *v)* Non logarithmic contributions to V_{eff} at a given loop order can alter the qualitative picture of the “tree-level improved” effective potential.

In this short presentation we concentrate on the interplay between *iv)* and *v)*, which leads, as we will see, to a modification of the usual discussion of EWSB conditions in the MSSM.

The 1-loop effective potential has the well-known form [3]

$$V = V_{tree} + \frac{\hbar}{64\pi^2} Str[M^4(Log \frac{M^2}{\mu_R^2} - 3/2)] \quad (1)$$

in the $\overline{MS}$ scheme. Here μ_R denotes the renormalization scale, V_{tree} the tree-level MSSM potential [4], M^2 the field dependent squared mass matrix of the scalar fields, and $Str[...] \equiv \sum_{spin} (-1)^{2s} (2s+1) (...)_s$, summing over gauge boson, fermion and scalar contributions. In the MSSM the explicit expression of $Str[M^4(Log \frac{M^2}{\mu_R^2} - 3/2)]$ is overwhelmingly lengthy. We have determined its exact structure analytically, relying on symbolic computation, thus allowing the exploration of virtually any direction in field space [5]. Hereafter we will be interested in the (H_1, H_2) manifold, *i.e.* all scalar fields but the two Higgs doublets are put to zero; $H_1 \equiv (H_1^0, H_1^-)$, $H_2 \equiv (H_2^+, H_2^0)$ consist of 8 real valued fields. Furthermore we limit the study here to $Str[M^4]$, that is we assume that logarithms in eq.(1) are reabsorbed in the running of all the parameters in $\bar{V}_{tree}(\mu_R^2)$ ². From this point of view $(-3/2)\hbar/64\pi^2 Str[M^4]$ is the genuine residual 1-loop effect. The effective potential takes then the form

$$V = \bar{V}_{tree}(\mu_R^2) + \frac{\hbar}{64\pi^2} (-3/2) Str M^4 \quad (2)$$

Now comes the main point: the usual EWSB conditions [1],

$$\frac{1}{2} M_Z^2 = \frac{\bar{m}_1^2 - \bar{m}_2^2 \tan^2 \beta}{\tan^2 \beta - 1} \quad , \quad \sin 2\beta = \frac{2\bar{m}_3^2}{\bar{m}_1^2 + \bar{m}_2^2} \quad (3)$$

² bypassing difficulties related to points *i)* and *iii)* above, as is often implicitly done in the literature! see however [6].

being merely the expressions of vanishing *first order* derivatives in the neutral directions $\langle H_1 \rangle = (v_1, 0)$, $\langle H_2 \rangle = (0, v_2)$, do not as such guarantee that we have indeed a (local) minimum of the effective potential at the electroweak scale, see point iv) above. However, provided that the $\overline{m}_i$'s are v_1 and v_2 independent, one can show that eq.(3), supplemented with the boundedness-from-below condition $\overline{m}_1^2 + \overline{m}_2^2 - 2|\overline{m}_3|^2 \geq 0$ does indeed imply that the 8 eigenvalues of the *second order* derivatives matrix are positive. It is noteworthy that this connection between 1st and 2nd order derivatives holds at tree-level, due to the special form of V_{tree} as dictated by softly broken supersymmetry as well as to the fact that the $\overline{m}_i$'s are all trivially v_1, v_2 independent at this level. It also continues to hold for the same reasons in the tree-level improved approximation ($\overline{V}_{tree}(\mu_R^2)$) where the running $\overline{m}_i(\mu_R^2)$ are v_1, v_2 independent as well ². [This is why eqs.(3) are often referred to in the literature as the *minimization conditions*, although they are not intrinsically so.] Nonetheless, there is no reason to believe that such a connection will hold for the full theory. Actually as we will see, it will already break down when non logarithmic 1-loop corrections are included, eq.(2). Indeed in this case, even though eqs.(3) retain their generic form, through a redefinition of the $\overline{m}_i$'s, there is now an explicit dependence on v_1 and v_2 in the $\overline{m}_i$'s. Injecting the full H_1, H_2 contribution of $StrM^4$ and extracting this explicit dependence, one finds the conditions for vanishing 1st order derivatives:

$$X_{m_1}^2 v_1 - X_{m_3}^2 v_2 + X v_1 (v_1^2 - v_2^2) + \tilde{\alpha} v_1^3 = 0 \quad , \quad X_{m_2}^2 v_2 - X_{m_3}^2 v_1 + X v_2 (v_1^2 - v_2^2) - \tilde{\alpha} v_2^3 = 0 \quad (4)$$

Here $X_{m_i} = m_i + \hbar \delta m_i$ and $X = g^2/8 + \hbar \delta g$, where δm_i (resp. δg) have been determined analytically and depend on all mass parameters of the MSSM and Yukawa couplings (resp. gauge couplings). $\tilde{\alpha} \equiv -9\hbar/(256\pi^2)g^2(Y_t^2 - Y_b^2)$ is a purely residual one-loop contribution. [$g^2 \equiv g_1^2 + g_2^2$, g_1 (resp. g_2) being the $U(1)$ (resp. $SU(2)_L$) gauge couplings, and we show in $\tilde{\alpha}$ just the top/bottom Yukawa dependence]. It is important to note at this level that a non zero $\tilde{\alpha}$ in eq.(4) modifies drastically the $\tan \beta$ dependence in the EWSB conditions. While these conditions are quadratic in $\tan \beta$ when $\tilde{\alpha} = 0$, (leading to eqs.(3) with v_1, v_2 free $\overline{m}_i$'s), they become quartic in $\tan \beta$ (and linear in $u \equiv v_1 v_2$) when $\tilde{\alpha} \neq 0$. More importantly now, one can show analytically that eqs.(4) together with the 1-loop boundedness-from-below condition $X_{m_1}^2 + X_{m_2}^2 - 2|X_{m_3}|^2 \geq 0$ no more suffice to characterize a local minimum and one has to resort explicitly to the positiveness conditions of all the physical Higgs squared masses³. However, working out the details [5], it turns out that only two independent new conditions are required and read when coupled with eqs.(4) (we give here for conciseness an approximate form of the first condition) :

$$a) \tan \beta \leq t_- \text{ or } \tan \beta \geq t_+ \quad (5)$$

$$\text{where } t_{\pm} \simeq 1 \pm \frac{4}{g} \sqrt{\frac{|\tilde{\alpha}|u}{X_{m_3}^2}} \sqrt{|\tilde{\alpha}|} + O(\tilde{\alpha}) \quad (6)$$

$$b) \frac{X_{m_3}^2}{u} \geq 0 \Leftrightarrow \tan \beta \geq T_+ \text{ (if } \tan \beta > 1) \text{ or } \tan \beta \geq T_- \text{ (if } \tan \beta < 1) \quad (7)$$

$$\text{where } T_{\pm} = \frac{X_{m_1}^2 + X_{m_2}^2 \pm \sqrt{(X_{m_1}^2 + X_{m_2}^2)^2 - 4X_{m_3}^4}}{2X_{m_3}^2} \quad (8)$$

³ Three of the eight eigenvalues are of course still vanishing and correspond to the goldstone components.

A couple of remarks are in order. Eqs.(4) and conditions *a)* and *b)* above, are necessary and sufficient conditions to 1-loop order for a local minimum to occur in the neutral $< H_1 > = (v_1, 0), < H_2 > = (0, v_2)$ direction. This is due to that on one hand a charge breaking vacuum does not occur in the H_1, H_2 sector (even to 1-loop order [5]), contrary to the non supersymmetric two-higgs-doublet case [8], and on the other hand, eqs.(4-8) imply by themselves that zero field configurations are energetically unstable, thus leading to spontaneous symmetry breaking. The boundary values $t_{\pm}$ in eq. (6) depend implicitly on $\tan\beta$ through u , the latter being determined by solving eqs.(4), where $\tilde{\alpha}u$ is formally $O(1)$ in $\tilde{\alpha}$. Furthermore, one has generically $T_+ > t_+$ and $T_- < t_-$. Conditions *a)* and *b)* exclude altogether a *calculable* region around $\tan\beta \sim 1$ in the otherwise unconstrained general MSSM. This should be contrasted with the usually quoted model-dependent constraint $1 \leq \tan\beta (\leq m_t/m_b)$ which requires an assumption of universality at the unification scale [7]⁴. It is also reassuring that in the limit $\hbar \rightarrow 0$, the values of $T_{\pm}$ degenerate into the $\tan\beta$ solutions for eqs.(3) while $t_{\pm} \rightarrow 1$, so that *a)* and *b)* are not constraints any more in this limit, as it should in the tree-level approximation.

Although valid in a model-independent context, it is obvious that conditions *a)* and *b)* can also be included consistently in a SUGRA-GUT motivated analysis. For instance in the 1-loop order approximation, one runs just the tree-level parts of the X_{m_i} 's and X , from say a unification scale downwards, while their 1-loop residual parts and $\tilde{\alpha}$ do not run. Eqs.(4) are then solved for $\tan\beta$ and u and the solutions not fulfilling eqs.(5-8) eliminated. It should be clear that this is done *prior* to any other physical requirement like reproducing the correct M_Z at the relevant scale, constraints on M_{top} , proton life-time, etc..., and as such could have a feedback on these requirements. Finally, even if the requirement of positive squared physical Higgs masses can be accounted for in a numerical analysis, we have established here analytically its theoretical connection with a model-independent constraint on the allowed values of $\tan\beta$.

This work was done in collaboration with Christophe Le Mouél. I would like to thank Gordon Kane, Dmitri Kazakov and Nikolai Krasnikov for suggestions and useful discussions during the conference.

References

- [1] For a pedagogical review, see for instance R. Arnowit and P. Nath, Lecture at Swieca School, Campos do Jordao, Brazil, Jan. 1993, and references therein
- [2] see for instance J.A. Casas, A. Lleyda, C. Muñoz, Nucl. Phys. B471 (1995) 3, and references therein;
- [3] S. Coleman, E. Weinberg, Phys. Rev. D 7 (1973) 1888;
- [4] For a review of the MSSM and related topics see H. P. Nilles, Phys. Rep. **110**(1984)1; H. E. Haber and G. L. Kane, Phys. Rep. **117**(1985)75; A. B. Lahanas and D. V. Nanopoulos, Phys. Rep. **145**(1987)1;
- [5] C. Le Mouél, G. Moulaka, in preparation;
- [6] C. Ford, D.R.T. Jones, P.W. Stephenson, M.B. Einhorn Nucl. Phys. B 395 (1993) 17; M. Bando, T. Kugo, N. Maekawa, H. Nakano, Phys. Lett. B 301 (1993) 83 and Prog.Theor.Phys.90 (1993) 405;
- [7] G.F. Giudice, G. Ridolfi, Z. Phys. C 41 (1988) 447;
- [8] M. Sher Phys. Rep. 179, Nos 5& 6 (1989) 273;

⁴ the other argument in favour of $1 \leq \tan\beta (\leq m_t/m_b)$, sometimes quoted in connection with perturbativity, is less quantitative and suffers from a less physical setting .